arXiv:1106.5202v3 [hep-ph] 8 Mar 2012

OU-HET-711/2011, MISC-2011-11

Neutrino Mass Matrix with No Adjustable Parameters

Yoshio Koide[a] **and Hiroyuki Nishiura**[b]

[a] *Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan*
E-mail address: koide@het.phys.sci.osaka-u.ac.jp

[b] *Faculty of Information Science and Technology, Osaka Institute of Technology, Hirakata, Osaka 573-0196, Japan*
E-mail address: nishiura@is.oit.ac.jp

Abstract

On the basis of the so-called "yukawaon" model, we found out a special form of the neutrino mass matrix M_ν which gives reasonable predictions. The M_ν is given by a multiplication form made of charged lepton mass matrix M_e and up-quark mass matrix M_u. This M_ν has no adjustable parameters except for those in M_e and M_u. Here, M_e and M_u are described by one parameter a_e (real) and two parameters a_u (complex), respectively, and those parameters are constrained by their observed mass ratios. With this form of M_ν, in spite of having only three parameters, the M_ν can give reasonable predictions $\sin^2 2\theta_{atm} \simeq 0.99$, $\sin^2 2\theta_{13} \simeq 0.015$, $\Delta m^2_{21}/\Delta m^2_{32} \simeq 0.030$, $\langle m_{ee}\rangle \simeq 0.0039$ eV, and so on, by using observed values of m_e/m_μ, m_μ/m_τ, m_c/m_t, and $\sin^2\theta_{solar}$ as input values.

PACS numbers: 14.60.Pq, 11.30.Hv, 12.60.-i,

1 Introduction

The observed masses and mixings of the quarks and leptons will provide a promising clue to a unified understanding of those fundamental particles. For such the purpose, not only investigating a theoretical model, but also searching for phenomenological mass matrix seem to be still effective. As one of phenomenological mass matrix models of the quarks and leptons, the so-called "yukawaon" model [1] (a kind of "flavon" models [2]) has been proposed. Here, the "effective" Yukawa coupling constants Y_f^{eff} are given by

$$(Y_f^{eff})_{ij} = \frac{y_f}{\Lambda}\langle (Y_f)_{ij}\rangle \qquad (i,j=1,2,3), \tag{1.1}$$

Λ is an energy scale of the effective theory, and $\langle Y_f\rangle$ are vacuum expectation value (VEV) matrices of scalar fields Y_f with 3×3 components. (Hereafter we call fields Y_f "yukawaons".) The most characteristic point in the yukawaon model is that all VEVs $\langle Y_f\rangle$ are described in terms of only one fundamental VEV matrix $\langle \Phi_e\rangle \propto \mathrm{diag}(\sqrt{m_e}, \sqrt{m_\mu}, \sqrt{m_\tau})$. For example, in a yukawaon model with an O(3) family symmetry [3, 4], the charged lepton, neutrino, up-quark, and down-quark mass matrices M_e, M_ν, M_u and M_d are given by

$$M_e \propto \langle Y_e \rangle \propto \langle \Phi_e \rangle \langle \Phi_e \rangle, \tag{1.2}$$

$$M_\nu = m_D M_R^{-1} m_D^T, \tag{1.3}$$

$$m_D \propto \langle Y_e \rangle, \quad M_R \propto \langle Y_R \rangle + m_{0\nu}^{-1} \langle Y_e \rangle \langle Y_e \rangle, \tag{1.4}$$

$$\langle Y_R \rangle \propto \langle \Phi_u \rangle \langle P_u \rangle \langle Y_e \rangle + \langle Y_e \rangle \langle P_u \rangle \langle \Phi_u \rangle + \xi_\nu \left(\langle \Phi_u \rangle \langle P_u \rangle \langle Y_e \rangle + \langle Y_e \rangle \langle P_u \rangle \langle \Phi_u \rangle \right). \tag{1.5}$$

$$M_u \propto \langle Y_u \rangle \propto \langle \Phi_u \rangle \langle \Phi_u \rangle, \tag{1.6}$$

$$\langle \Phi_u \rangle \propto \langle \Phi_e \rangle (\langle E \rangle + a_u \langle X \rangle) \langle \Phi_e \rangle, \tag{1.7}$$

$$M_d \propto \langle Y_d \rangle \propto \langle \Phi_e \rangle (\langle E \rangle + a_d \langle X \rangle) \langle \Phi_e \rangle, \tag{1.8}$$

where

$$\langle E \rangle = v_E \mathbf{1} = v_E \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \langle X \rangle = v_X S_3 \equiv \frac{1}{3} v_X \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \tag{1.9}$$

in the diagonal basis of M_e, while $\langle P_u \rangle$ is given by a form

$$\langle P_u \rangle_u \propto \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{1.10}$$

in the diagonal basis of M_u.

In this paper, we will find out a special form of the neutrino mass matrix which is compatible with the observed neutrino data in spite of having no adjustable parameters. The form will be obtained along the lines of the yukawaon model by changing the structure of $\langle Y_e \rangle$ from Eq.(1.2).

The neutrino mass matrix is still given by the form of seesaw type, $M_\nu = m_D M_R^{-1} m_D^T$ with $m_D \propto \langle Y_e \rangle$, but the Majorana neutrino mass matrix M_R is changed into a simple form

$$M_R \propto \langle Y_R \rangle \propto \langle \Phi_u \rangle \langle Y_e \rangle + \langle Y_e \rangle \langle \Phi_u \rangle. \tag{1.11}$$

Here, we assume that the charged lepton mass matrix M_e is given by

$$\langle Y_e \rangle \propto \langle \bar{\Phi}_0 \rangle (\langle E' \rangle + a_e \langle X_2 \rangle) \langle \bar{\Phi}_0 \rangle, \tag{1.12}$$

differently from Eq.(1.2), and we also redefine $\langle \Phi_u \rangle$ as

$$\langle \Phi_u \rangle \propto \langle \Phi_0 \rangle (\langle E \rangle + a_u \langle X_3 \rangle) \langle \Phi_0 \rangle, \tag{1.13}$$

where $\langle E' \rangle = \langle E \rangle = v_E \mathbf{1}$ and

$$\langle X_2 \rangle = v_X S_2 \equiv \frac{1}{2} v_X \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \langle X_3 \rangle = v_X S_3 \equiv \frac{1}{3} v_X \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \tag{1.14}$$

Note that there are no ξ_ν term, no $m_{0\nu}^{-1}$ term and no $\langle P_u \rangle$ in Eq.(1.11), i.e. M_ν is simply given by

$$M_\nu \simeq k_\nu \left(M_e^{-1} M_u^{1/2} + M_u^{1/2} M_e^{-1} \right). \tag{1.15}$$

In other words, there is no adjustable parameter in the present neutrino mass matrix, except for a_e in M_e and a_u in M_u^{12} (i.e. $\langle \Phi_u \rangle$). The purpose of the present paper is not to derive the mass matrix forms (1.11) - (1.13) theoretically, but to demonstrate that the phenomenological neutrino mass matrix (1.15) with Eqs.(1.11) - (1.13) can be compatible with the present neutrino data in spite of quite few parameters. We predict $\sin^2 2\theta_{atm} \simeq 0.99$, $\sin^2 2\theta_{13} \simeq 0.015$, $\Delta m_{21}^2 / \Delta m_{32}^2 \simeq 0.030$, $\langle m_{ee} \rangle \simeq 0.0039$ eV, and so on, by using observed values of m_e/m_μ, m_μ/m_τ, and $\sin^2 \theta_{solar}$ as input values. In this paper, for simplicity, we do not discuss the down-quark mass matrix M_d and the Cabibbo-Maskawa- Kobayashi [5] (CKM) mixing.

In the next section, superpotentials for yukawaons and the assignments of the fields in the present U(3) yukawaon model are investigated. In Sec. 3, numerical results of the model are discussed. Sec. 4 is devoted to the concluding remarks. In Appendix A, R-charge assignments are discussed. In Appendix B, we present a rotation matrix which transforms $\langle X_3 \rangle$ into $\langle X_2 \rangle$.

2 Superpotential

In this section, we give superpotentials for the yukawaons and the assignments of the fields in the present U(3) yukawaon model.

In the yukawaon model, the order of the fields is important. Therefore, in this paper, let us assume a U(3) family symmetry instead of O(3) and denote fields $\mathbf{6}^*$ and $\mathbf{6}$ of U(3) as $\bar{A}$ and A, respectively. (Therefore, it should be noted that a term $\bar{A}B\bar{C}$ is allowed, but $\bar{A}\bar{C}B$ and $B\bar{A}\bar{C}$ are forbidden.) In the U(3) model, for example, the relation (1.6) is re-expressed as

$$(M_u)^{ij} \propto \langle \bar{Y}_u^{ij} \rangle \propto \langle \bar{\Phi}_u^{ik} \rangle \langle E_{kl}^u \rangle \langle \bar{\Phi}_u^{lj} \rangle, \tag{2.1}$$

with $\langle E^u \rangle = v_E \mathbf{1}$. In order to distinguish each yukawaon from other yukawaons, although we assumed $\mathrm{U}(1)_X$ charge in the O(3) model [3, 4], in this U(3) model, we assume only R charge conservation instead of $\mathrm{U}(1)_X$ charge conservation. For the right handed neutrino sector ($\bar{Y}_R$), it should be noted that we cannot add $\bar{Y}_e \bar{Y}_e$ term to $\bar{Y}_R$ as in Eq.(1.4). In the old model, we assigned the $\mathrm{U}(1)_X$ charges Q_X only for gauge singlet fields, e.g. $Q_X(\ell) = Q_X(H_d)$, i.e. $Q_X(Y_e) = -Q(e^c)$. Besides, we assumed $Q_X(e^c) = Q(\nu^c)$ in order to build a model without Y_ν. Therefore, we could obtain $Q_X(Y_R) = Q_X(Y_e Y_e)$ in the old model. However, in this U(3) model, we cannot obtain $R(Y_R) = R(\bar{Y}_e E \bar{Y}_e)$. Besides, we cannot introduce a ξ_ν term such as in Eq.(1.5).

We assume the following superpotential $W = W_Y + W_e + W_d + W_u' + W_u + W_R + W_E$:

$$W_Y = \frac{y_e}{\Lambda} \ell_i \bar{Y}_e^{ij} e_j^c H_d + \frac{y_\nu}{\Lambda} \ell_i \bar{Y}_e^{ij} \nu_j^c H_u + \lambda_R \nu_i^c \bar{Y}_R^{ij} \nu_j^c$$

$$+ \frac{y_u}{\Lambda} u_i^c \bar{Y}_u^{ij} q_j H_u + \frac{y_d}{\Lambda} d_i^c \bar{Y}_d^{ij} q_j H_d + \mu_H H_u H_d, \tag{2.2}$$

	H_u	H_d	E^u	$\bar{E}_u$	Θ_{8+1}
U(3)	**1**	**1**	**6**	**6***	**8 + 1**
R	1	1	$1-\bar{r}_E$	$\bar{r}_E$	1
Model	1	1	3	-2	1

ℓ	e^c	ν^c	$\bar{Y}_e$	$\bar{\Phi}_0$	E'	X'	Θ^e	$\bar{Y}_R$	Θ^R
3	**3**	**3**	**6***	**6***	**6**	**6**	**6**	**6***	**6**
r_ℓ	r_e	r_e	r_{Ye}	r_0	$r_{X'}$	$r_{X'}$	$2-r_{Ye}$	r_R	$2-r_R$
1	0	0	0	$\frac{1}{2}$	-1	-1	2	2	0

q	u^c	d^c	$\bar{Y}_u$	$\bar{\Phi}_u$	Θ^u	$\Theta^{u\prime}$	$\bar{Y}_d$	Θ^d	E	X
3	**3**	**3**	**6***	**6***	**6**	**6**	**6***	**6**	**6**	**6**
r_q	r_u	r_d	r_{Yu}	r_{Yd}	$2-r_{Yu}$	$2-r_{Yd}$	r_{Yd}	$2-r_{Yd}$	r_X	r_X
1	-1	$+1$	$+1$	-1	1	3	-1	3	-2	-2

Table 1: Assignments of R charges, where $r_{X'} = r_{Ye} - 2r_0$, $r_X = r_{Yd} - 2r_0$ and $R(\bar{\Phi}u) = R(\bar{Y}_d) \equiv r_{Yd}$. The values in the third raw denote R charge values in a special case under the assumptions (A.11) and (A.14). For more details, see Eqs.(A.1) - (A.20) in Appendix A.

$$W_e = \mu_e \mathrm{Tr}[\bar{Y}_e \Theta^e] + \frac{\lambda_e}{\Lambda}\mathrm{Tr}[\bar{\Phi}_0(E' + a_e X_2)\bar{\Phi}_0\Theta^e], \tag{2.3}$$

$$W_d = \mu_d \mathrm{Tr}[\bar{Y}_d \Theta^d] + \frac{\lambda_d}{\Lambda}\mathrm{Tr}[\bar{\Phi}_0(E + a_d e^{i\alpha_d} X_3)\bar{\Phi}_0\Theta^d], \tag{2.4}$$

$$W'_u = \mu'_u \mathrm{Tr}[\bar{\Phi}_u \Theta^{u\prime}] + \frac{\lambda'_u}{\Lambda}\mathrm{Tr}[\bar{\Phi}_0(E + a_u e^{i\alpha_u} X_3)\bar{\Phi}_0\Theta^{u\prime}], \tag{2.5}$$

$$W_u = \mu_u \mathrm{Tr}[\bar{Y}_u \Theta^u] + \frac{\lambda_u}{\Lambda}\mathrm{Tr}[\bar{\Phi}_u E^u \bar{\Phi}_u \Theta_u] + \frac{\lambda_{0u}}{\Lambda^5}\mathrm{Tr}[(\bar{E}_u E^u)^3 \bar{E}_u \Theta^u], \tag{2.6}$$

$$W_R = \mu_R \mathrm{Tr}[\bar{Y}_R \Theta^R] + \frac{\lambda_R}{\Lambda}\mathrm{Tr}[\left(\bar{\Phi}_u E^u \bar{Y}_e + \bar{Y}_e E^u \bar{\Phi}_u\right)\Theta^R], \tag{2.7}$$

where, in Eq.(2.2), q and ℓ are $\mathrm{SU}(2)_L$ doublet fields, and f^c ($f = u, d, e, \nu$) are $\mathrm{SU}(2)_L$ singlet fields. The other fields in Eqs.(2.3)-(2.7) have quantum numbers defined in Table 1.

In Eq.(2.6), the third term has been added since it has the same R charge as that of $\bar{Y}_u\Theta^u$. (See Eq.(A.20) in Appendix A.) The λ_{0u} term plays a role in shifting eigenvalues of the up-quark mass matrix M_u by a constant value. Details of R charge assignments are given in Appendix A.

For the field E^u, we assume an additional field $\bar{E}_u$, and consider a superpotential with a form

$$W_E = \lambda_E \mathrm{Tr}[E^u \bar{E}_u \Theta_{8+1}] + \lambda'_E \mathrm{Tr}[E^u \bar{E}_u]\,\mathrm{Tr}[\Theta_{8+1}], \tag{2.8}$$

where Θ_{8+1} is a field $\mathbf{8}+\mathbf{1}$ of U(3) with $\langle\Theta_{8+1}\rangle = 0$. The superpotential W_E leads to $\langle E^u\rangle\langle\bar{E}_u\rangle \propto \mathbf{1}$. We assume that the form

$$\langle\bar{E}_u\rangle \propto \langle E^u\rangle = v_E\,\mathrm{diag}(1,1,1), \tag{2.9}$$

is given by a specific form of the solutions $\langle E^u\rangle\langle\bar{E}_u\rangle \propto \mathbf{1}$.

In this paper, we do not discuss a superpotential which gives the observed charged lepton mass spectrum. We only use the observed charged lepton mass values as input values in $\langle\bar{Y}_e\rangle_e$.

Under the assumption that all Θ fields take $\langle\Theta\rangle = 0$, SUSY vacuum conditions lead to VEV relations[1] from Eqs.(2.3)-(2.7). That is, instead of Eqs.(1.2) - (1.8) in the previous model, we obtain the following mass matrix relations:

$$M_e \propto \langle\bar{Y}_e\rangle \propto \langle\bar{\Phi}_0\rangle(\mathbf{1} + a_e S_2)\langle\bar{\Phi}_0\rangle, \tag{2.10}$$

$$M_d \propto \langle\bar{Y}_d\rangle \propto \langle\bar{\Phi}_0\rangle(\mathbf{1} + a_d e^{i\alpha_d} S_3)\langle\bar{\Phi}_0\rangle, \tag{2.11}$$

$$\langle\bar{\Phi}_u\rangle \propto \langle\bar{\Phi}_0\rangle(\mathbf{1} + a_u e^{i\alpha_u} S_3)\langle\bar{\Phi}_0\rangle, \tag{2.12}$$

$$M_u \propto \langle\bar{Y}_u\rangle \propto \langle\bar{\Phi}_u\rangle\cdot\mathbf{1}\cdot\langle\bar{\Phi}_u\rangle + (v_{\Phi u})^2\zeta_u\mathbf{1}, \tag{2.13}$$

$$M_R \propto \langle\bar{Y}_R\rangle \propto \langle\bar{\Phi}_u\rangle\cdot\mathbf{1}\cdot\langle\bar{Y}_e\rangle + \langle\bar{Y}_e\rangle\cdot\mathbf{1}\cdot\langle\bar{\Phi}_u\rangle. \tag{2.14}$$

Here, we can take a diagonal basis of $\langle\bar{\Phi}_0\rangle$ without loosing the generality:

$$\langle\bar{\Phi}_0\rangle = \mathrm{diag}(v_1, v_2, v_3) = v_0\,\mathrm{diag}(x_1, x_2, x_3), \tag{2.15}$$

where we have normalized x_i as $x_1^2 + x_2^2 + x_3^2 = 1$. The neutrino mass matrix M_ν is given by a seesaw type $M_\nu = m_D M_R^{-1} m_D^T$ with $m_D \propto M_e$ similar to Eq.(1.4) [but there is no $m_{0\nu}^{-1}$]. Note that the previous relations (1.2) - (1.8) were given at a diagonal basis of the VEV $\langle\Phi_e\rangle$, while present relations (2.10) - (2.14) are given at a diagonal basis of the VEV $\langle\bar{\Phi}_0\rangle$. Here the numerical matrices S_3 and S_2 are defined by

$$S_3 = \frac{1}{3}\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad S_2 = \frac{1}{2}\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{2.16}$$

at the diagonal basis of the VEV $\langle\bar{\Phi}_0\rangle$. Note that the VEV matrix $\langle\bar{Y}_e\rangle$ in Eq.(2.10) is no more diagonal in this basis.

In obtaining the mixing matrices, the common coefficients are not important. Here we have taken $v_{E'} = v_{X2}$ and $v_E = v_{X3}$ for simplicity. The ζ_u term in Eq.(2.13) comes from the new term given in Eq.(A.20). This term contributes to the up-quark mass ratios, while not to the up-quark mixing matrix, so that it does not change the predictions for the neutrino mixing parameters. We suppose that the contribution from such the higher dimensional term (A.20) is considerably small, so that it also does not visibly affect the up-quark mass ratio m_c/m_t, although it can slightly affect m_u/m_c.

[1] For example, in obtaining the relation (2.10), we have assumed a vacuum with $\langle\Theta^e\rangle = 0$, so that the conditions $\partial W/\partial Y_e = 0$ and $\partial W/\partial\Phi_0 = 0$ do not affect other VEV relations obtained from SUSY vacuum conditions $\partial W/\partial\Theta_A = 0$ ($A \neq e$). We assume that the observed SUSY symmetry breaking is induced by a gauge mediation mechanism (not including family symmetry), so that our VEV relations among yukawaons are still valid in the quark and lepton sectors after the SUSY is broken .

3 Numerical results in the up-quark and neutrino mass matrices

In this section, we investigate whether the new VEV matrix relations (2.10) - (2.14) can well describe the observed neutrino mixing parameters together with the observed up-quark mass ratios or not.

Since the charged lepton mass matrix given by Eq.(2.10) is not diagonal, the lepton mixing matrix [Pontecorvo-Maki-Nakagawa-Sakata (PMNS) [7] mixing matrix] U in the present conventions is defined by

$$U = U_{eL}^{\dagger} U_{\nu L}, \tag{3.1}$$

where U_{eL} and $U_{\nu L}$ are defined by

$$U_{eL}^{T} \langle \bar{Y}_e \rangle U_{eL} = \langle \bar{Y}_e \rangle^{diag} \equiv \langle \bar{Y}_e \rangle_e, \tag{3.2}$$

$$U_{\nu L}^{\dagger} (M_\nu^{\dagger} M_\nu) U_{\nu L} = (M_\nu^{\dagger} M_\nu)^{diag}, \tag{3.3}$$

and M_ν is given by

$$M_\nu = \frac{y_\nu^2}{\lambda_R} \left(\frac{\langle H_u^0 \rangle}{\Lambda} \right)^2 \langle \bar{Y}_e \rangle \langle \bar{Y}_R \rangle^{-1} \langle \bar{Y}_e \rangle. \tag{3.4}$$

Neutrino mixing parameters we discuss are $\tan^2 \theta_{solar} = |U_{12}|^2/|U_{11}|^2$, $\sin^2 2\theta_{atm} = 4|U_{23}|^2|U_{33}|^2$, and $|U_{13}|^2$. Here U_{ij} are the matrix elements of the lepton mixing matrix defined by (3.1).

The matrix M_u in (2.13) is diagonalized as

$$U_{uL}^{\dagger} (M_u^{\dagger} M_u) U_{uL} = (M_u^{\dagger} M_u)^{diag}. \tag{3.5}$$

Here U_{uL} is a mixing matrix among left-handed up-quarks u_{Li} . (In the present paper, the mass matrices (i.e. $\langle \bar{Y}_f \rangle$) are defined by Eq.(2.2). Therefore, the conventions of the mixing matrices are somewhat changed from the conventional ones.) Note that since the VEV matrix $\langle \bar{\Phi}_u \rangle$ is complex and $\langle \bar{Y}_u \rangle$ is given by Eq.(2.13), the diagonalization of the up-quark mass matrix must be done by Eq.(3.5).

3.1 Parameters in the model

The mass matrices for quarks and neutrinos in the O(3) model have been described in terms of the fundamental VEV matrix $\langle \Phi_e \rangle$. On the other hand, the fundamental VEV matrix in the present model is $\langle \bar{\Phi}_0 \rangle$ defined by Eq.(2.10) in which we have new parameter a_e. Thus the number of parameters are increased by one compared with the previous model (1.2). On the other hand, we cannot bring neither the ξ_ν term given in Eq.(1.5) nor $\langle P_u \rangle_u$ defined in Eq.(1.10) into the present model, so that there are no parameters which are corresponding to ξ_ν and P_u.

The VEV of $\langle \bar{\Phi}_0 \rangle = \mathrm{diag}(v_1, v_2, v_3)$ is related to the charged lepton mass matrix M_e as follows:

$$M_e = k \begin{pmatrix} \left(1 + \frac{1}{2} a_e\right) v_1^2 & \frac{1}{2} a_e v_1 v_2 & 0 \\ \frac{1}{2} a_e v_1 v_2 & \left(1 + \frac{1}{2} a_e\right) v_2^2 & 0 \\ 0 & 0 & v_3^2 \end{pmatrix}, \tag{3.6}$$

where $k = -(\lambda_e/\mu_e\Lambda)(y_e\langle H_d^0\rangle/\Lambda)v_{X'}$, so that we obtain

$$m_e + m_\mu = k\left(1 + \frac{1}{2}a_e\right)(v_1^2 + v_2^2), \tag{3.7}$$

$$m_e m_\mu = k^2(1 + a_e)v_1^2 v_2^2, \tag{3.8}$$

and $m_\tau = kv_3^2$. Here, since we are interested only in the relative ratios among the eigenvalues of the charged lepton mass matrix M_e, the common coefficient k is not a parameter of the model. The 3 parameters a_e, v_1/v_2 and v_2/v_3 are sufficient to determine the two charged lepton mass ratios and charged lepton mixing matrix U_e which is described only by one parameter θ_{12}^e. [The mixing angle θ_{12}^e is not observable. The observed quantities are parameters of the lepton mixing matrix defined by Eq.(3.1).] Therefore, when we give a value of the parameter a_e, the values of v_i are completely determined by the input values of the charged lepton masses. In other words, even when we give three charged lepton masses as the inputs, one of the free parameters still remains.

Thus, in the present model, we have 4 parameters a_e, a_u, α_u and ζ_u (except for the input values m_e, m_μ, and m_τ) for the up-quark and neutrino mass matrices. On the other hand, the number of the predictable quantities are 12, i.e., 2+2+2 mass ratios (up-quark, charged lepton and neutrino mass ratios) and 4+2 PMNS mixing parameters (including two Majorana phases). At present, we know 6 observed values of $\sqrt{m_u/m_c}$, $\sqrt{m_c/m_t}$, $\tan^2\theta_{solar}$, $\sin^2 2\theta_{atm}$, $\sin^2 2\theta_{13}$, and $R_\nu = \Delta m_{21}^2/\Delta m_{32}^2$ in addition to the charged lepton masses.

The term $\zeta_u \mathbf{1}$ in M_u given in Eq.(2.13) does not affect the up-quark mixing matrix, so that it also affects neither quark or lepton mixing matrices. Since we suppose $|\zeta_u|^2 \ll 1$, the term almost does not affect $\sqrt{m_c/m_t}$, although it can slightly affect $\sqrt{m_u/m_c}$. As a result, the present model predicts 11 observables by using the three parameters (a_e, a_u, α_u). In other words, the value of $\sqrt{m_u/m_c}$ is not " prediction", and it is a quantity which can be adjustable by the additional parameter ζ_u freely.

3.2 Numerical results

Now let us show the results of numerical analysis of the model. First, we show, in Fig. 1, the a_u dependences of the quantities $\sqrt{m_c/m_t}$, $\tan^2\theta_{solar}$, and $\sin^2 2\theta_{atm}$ with taking typical values of $a_e = 3,\ 30,\ 100$ and $\alpha_u = 0^\circ,\ 15^\circ$ in order to see rough parameter behaviors.

As seen in Fig. 1, we can find that (i) the value of $\sqrt{m_c/m_t}$ takes a maximum value at $a_u \sim -3$ insensitively to the values of a_e and α_u; (ii) since the maximum value of $\sin^2 2\theta_{atm}$ shows $\sin^2 2\theta_{atm} \simeq 1$ which is in favor of the observed value, we must search for a parameter set (a_e, a_u, α_u) which gives a maximum value of $\sin^2 2\theta_{atm}$; (iii) a case with a small value of a_e gives a large value of $\tan^2\theta_{solar}$ compared with the observed value $\tan^2\theta_{solar} \sim 0.5$ (see Fig. 1 (a)), so that such a case is ruled out; on the other hand, a case with a large value of a_e gives a small $\tan^2\theta_{solar}$ (see Fig. 1 (c)), so that such a case is also ruled out; (iv) as a result, a region of (a_e, a_u) which can give $\sin^2 2\theta_{atm} \simeq 1$ and $\tan^2\theta_{solar} \sim 0.5$ is $(a_e, a_u) \sim (30, -3)$.

Next, in order to determine parameter values (a_e, a_u, α_u), let us illustrate, in Fig. 2, the α_u behaviors of $\sqrt{m_c/m_t}$, $\tan^2\theta_{solar}$ and $\sin^2 2\theta_{atm}$ at $a_e \sim 28$ and $a_u \sim -3$. From Fig. 2, we



Figure 1: $\sqrt{m_c/m_t}$, $\tan^2\theta_{solar}$, and $\sin^2 2\theta_{atm}$ versus a parameter a_u for typical parameter values $a_e = 3$ (Fig. 1 (a)), $a_e = 30$ (Fig. 1 (b)), and $a_e = 100$ (Fig. 1 (c)) with $\alpha_u = 0°$ (solid curves) and $\alpha_u = 15°$ (dashed curves). Curves "r23", "solar", and "atm" denote "r23"$= \sqrt{m_c/m_t} \times 10$, "solar"$= \tan^2\theta_{solar}$, and "atm"$= \sin^2 2\theta_{atm}$, respectively.



Figure 2: $\sqrt{m_c/m_t}$, $\tan^2\theta_{solar}$, and $\sin^2 2\theta_{atm}$ versus a parameter α_u for typical parameter values $a_e = 26$ (Fig. 2 (a)), $a_e = 28$ (Fig. 2 (b)), and $a_e = 30$ (Fig. 2 (c)) with $a_u = -2.9$ (dashed curves), $a_u = -3.0$ (solid curves), and $a_u = -3.1$ (dot-dashed curves). Curves "r23", "solar", and "atm" denote "r23"$= \sqrt{m_c/m_t} \times 10$, "solar"$= \tan^2\theta_{solar}$, and "atm"$= \sin^2 2\theta_{atm}$, respectively.

search for the value α_u which gives the observed value [8] $\sqrt{m_c/m_t} = 0.0600^{+0.0045}_{-0.0047}$ at $\mu = m_Z$. We find that the value $\alpha_u \simeq 15°$ can give a reasonable fit $\sqrt{m_c/m_t} = 0.0600$ insensitively to the other parameters.

Therefore, by fixing the value $\alpha_u = 15°$, we illustrate the contour lines of $\sqrt{m_c/m_t}$ and $\tan^2\theta_{solar}$ in the (a_e, a_u) plane in Fig. 3. The curves denote (a_e, a_u) which gives the observed values $\sqrt{m_c/m_t} = 0.0600^{+0.0045}_{-0.0047}$ [8] and $\tan^2\theta^{obs}_{solar} = 0.47^{+0.05}_{-0.03}$ [9]. As seen in Fig. 3, we have two intersection points of the curves of $\sqrt{m_c/m_t}$ and $\tan^2\theta_{solar}$. For the center values $\sqrt{m_c/m_t} = 0.060$ and $\tan^2\theta_{solar} = 0.47$, the solutions (a_e, a_u, α_u) are

$$(26.7, -2.88, 15°), \qquad (27.9, -3.18, 15°). \tag{3.9}$$



Figure 3: Contour lines of $\sqrt{m_c/m_t}$ and $\tan^2\theta_{solar}$ in the (a_e, a_u) plane in the case of $\alpha_u = 15°$. As input values, $\sqrt{m_c/m_t} = 0.0600^{+0.0045}_{-0.0047}$ and $\tan^2\theta^{obs}_{solar} = 0.47^{+0.05}_{-0.03}$ have been used. Curves with dot-dash, solid, and dash denote the upper, center, and lower observed values, respectively. Curves "r23" and "solar" denote "r23"$= \sqrt{m_c/m_t} \times 10$ and "solar"$= \tan^2\theta_{solar}$, respectively.

Input (a_e, a_u, α_u)	$\sqrt{m_c/m_t}$	$\tan^2\theta_{solar}$	$\sin^2 2\theta_{atm}$	$\sin^2 2\theta_{13}$	R_ν
$(26.7, -2.88, 15°)$	0.0603	0.470	0.990	0.015	0.0303
$(27.9, -3.18, 15°)$	0.0601	0.469	0.980	0.011	0.0204
upper	+0.0045	+0.05			+0.0027
Observed value	0.0600	0.47	> 0.92 [9]	< 0.15 [9]	0.0312
lower	−0.0047	−0.03			−0.0023

Table 2: Predicted values for the parameter values (a_e, a_u, α_u).

We list our prediction values for these parameter solutions in Table 1. Of the two solutions obtained from the input data $\sqrt{m_c/m_t}$ and $\tan^2\theta_{solar}$, Table 2 suggests that we should take the former one considering the observed value of R_ν [9]

$$R_\nu \equiv \frac{\Delta m^2_{solar}}{\Delta m^2_{atm}} = (3.12^{+0.27}_{-0.23}) \times 10^{-2}. \tag{3.10}$$

For reference, we also illustrate the behavior of predicted values for input values (a_e, a_u, α_u) around the parameter solutions (3.9) in Fig. 4. As seen in Fig. 4, the predicted values $\sin^2 2\theta_{atm}$ and $\tan^2\theta_{solar}$ are insensitive to the parameter values a_e, a_u, and α_u around the values $(a_e, a_u, \alpha_u) = (26.7, -2.88, 15°)$. However, $\sqrt{m_c/m_t}$ and $|U_{13}|^2$ (and also R_ν) are somewhat dependent on these parameters. Since these parameter values are mainly obtained by taking the input value $\sqrt{m_c/m_t} = 0.0600$, if the input value changes, then the predicted values will also change.

So far, we have not discussed the value of m_u/m_c. In the present model, the value of m_u/m_c is always adjustable by the parameter ζ_u given in Eq.(2.13) without affecting other



Figure 4: Predicted values versus a parameter α_u for $a_e = 26.7$ and $a_u = -2.88$. Curves "r23", "solar", "atm", "u13", and "R" denote "r23"$= \sqrt{m_c/m_t} \times 10$, "solar"$= \tan^2\theta_{solar}$, "atm"$= \sin^2 2\theta_{atm}$, "u13"$= |U_{13}|^2 \times 100$, and "R"$= R_\nu \times 10$, respectively. For reference, curves for $(a_e, a_u) = (26.7, -3.00)$ (dash curve), $(26.7, -2.80)$ (dot curve), $(29, -2.88)$ (dot dash curve), and $(25, -2.88)$ (2-dot dash curve) are illustrated in addition to the curve (solid) for $(26.7, -2.88)$.

Input ζ_u	$\sqrt{m_u/m_c}$	$\sqrt{m_c/m_t}$	$\tan^2\theta_{solar}$	$\sin^2 2\theta_{atm}$	$\sin^2 2\theta_{13}$	δ_{CP} (J)	R_ν
0	0.0180	0.0603	0.470	0.990	0.015	-102.5° (-0.0139)	0.0303
3.8×10^{-7}	0.0454	0.0602	0.470	0.990	0.015	-102.5° (-0.0139)	0.0303

Table 3: Predicted values versus ζ_u parameter. Other parameters (a_e, a_u, α_u) have taken the same values $(26.7, -2.88, 15^\circ)$ as those in Table 2.

predicted values. In order to fit the predicted value of $\sqrt{m_u/m_c}$ to the observed value [8] $\sqrt{m_u/m_c} = 0.0453^{+0.012}_{-0.010}$, we choose ζ_u as $\zeta_u = 3.8 \times 10^{-7}$. As seen in Table 3, the value of ζ_u almost does not change the numerical predictions given in Table 2.

In conclusion, we take the parameter set

$$(a_e, a_u, \alpha_u, \zeta_u) = (26.7, -2.88, 15^o, 3.8 \times 10^{-7}). \tag{3.11}$$

Then, we predict neutrino masses

$$m_{\nu 1} \simeq 0.012 \text{ eV}, \quad m_{\nu 2} \simeq 0.015 \text{ eV}, \quad m_{\nu 3} \simeq 0.051 \text{ eV}, \tag{3.12}$$

by using the input value $\Delta m^2_{32} \simeq 0.0024$ eV^2. We also predict the effective Majorana mass $\langle m_{ee} \rangle$ in the neutrinoless double beta decay [10]

$$\langle m_{ee} \rangle = \left| m_1 U^2_{e1} + m_2 U^2_{e2} + m_3 U^2_{e3} \right| \simeq 0.0039 \text{ eV}. \tag{3.13}$$

It is worthwhile noticing approximately degenerate neutrino masses $m_{\nu 1} \sim m_{\nu 2}$.

4 Concluding remarks

In this paper we have found out a special form of the neutrino mass matrix M_ν based on a yukawaon model with U(3) family symmetry, which has quite few free parameters. With this form of M_ν, the M_ν can give reasonable predictions in spite of having no adjustable parameters, i.e. the M_ν is simply given by the form (1.15), and the mass matrices M_e and M_u include parameters a_e and a_u, respectively, which are fixed by their observed mass ratios. In this yukawaon model, the yukawaon VEV matrices are described in terms of a new fundamental VEV matrix $\langle\bar{\Phi}_0\rangle$. For example, the yukawaon VEV matrix $\langle\bar{Y}_e\rangle$ for the charged leptons is given by (2.10) which has the structure of $(\mathbf{1}+a_eS_2)$ with a new parameter a_e. This structure in $\langle\bar{Y}_e\rangle$ has been chosen from a phenomenological point of view and there is no reason why $\langle\bar{Y}_e\rangle$ takes such a form. Nevertheless if we accept the form (2.10), then we can obtain a simple form of VEV matrix $\langle\bar{Y}_R\rangle$ for right-handed neutrinos without introducing the somewhat strange VEV matrix P_u and ξ_ν term that were introduced in the O(3) model to get the observed nearly tribimaximal neutrino mixing [11].

The new model has only four parameters $(a_e, a_u, \alpha_u, \zeta_u)$ as far as the up-quark and lepton sectors are concerned. on the other hand, we have 12 observable quantities (2 up-quark mass ratios, 4 lepton mass ratios, and 4+2 lepton mixing parameters). The parameter ζ_u affects only the prediction of m_u/m_c, so that we have fixed it by the observed value of $r^u_{12} = \sqrt{m_u/m_c}$. The parameter α_u is sensitive only to m_c/m_t, so that we have fixed by the observed value of $r^u_{23} = \sqrt{m_c/m_t}$ as seen in Fig. 2 (b). The parameter a_e is determined from the cross point of the predicted values of r^u_{23} and $\tan^2\theta_{solar}$ in the a_e-a_u plane. Note that we have used only the observed values of r^u_{23} and $\tan^2\theta_{solar}$ in order to fix the three parameters (a_e, a_u, α_u). Although we have tacitly used $\sin^2 2\theta_{atm} \sim 1$, we have not used the observed value of $\sin^2 2\theta_{atm}$ explicitly.

On the other hand, for the remaining 2 down-quark mass ratios and 4 CKM mixing parameters, we have additional 2 parameters (a_d and α_d). Regrettably, we cannot obtain reasonable predictions with the two parameters, although we can fit the values of down-quark mass ratios and V_{us}. The situation is the same as in the previous O(3) model. We must introduce a phase matrix P_d with two parameters (ϕ_1, ϕ_2) and a common mass shift term $m_{0d}\mathbf{1}$. Then, five parameters can fit six observables barely. Therefore, the model is not so attractive for down-quark sector. In this paper, we did not demonstrate the explicit numerical fitting for down-quark mass rations and CKM mixing parameters.

The present U(3) model have the following interesting features in the lepton sector:
(i) The model predicts $\sin^2 2\theta_{13} \sim 0.015$. In the previous O(3) model, the predicted value of $|U_{13}|^2$ was invisibly small, i.e. $|U_{13}|^2 \sim 10^{-4}$. The T2K experiment [12] put a constraint $0.03 < \sin^2 2\theta_{13} < 0.28$ (90% C.L.) for $\delta_{CP} = 0$ and a normal hierarchy. Our predicted value $\sin^2 2\theta_{13} = 0.015$ seems to be somewhat lower than the experimental lower bound. However, as seen in Table 3, our prediction on δ_{CP} gives $\delta_{CP} = -103^\circ$, which decreases the lower bound 0.03 of the T2K result to 0.02. Besides, the Double CHOOZ experiment [13] has reported that $\sin^2 2\theta_{13} = 0.085 \pm 0.029 \pm 0.042$ at 68% CL. The lower value is $\sin^2 2\theta_{13} = 0.014$. Therefore, we consider that the predicted value $\sin^2 2\theta_{13} = 0.015$ is yet not ruled out, although the status is considerably severe.

(ii) It also predicts a reasonable value of $R_\nu \equiv \Delta m^2_{solar}/\Delta m^2_{atm} \sim 0.03$ in contrast to the case of the O(3) model in which we could not predict R_ν. (In the previous model, the value of R_ν needed to adjust the additional free parameter $m_{0\nu}^{-1}$ in Eq.(1.4).)
(iii) The present model gives approximately degenerate neutrino masses $m_{\nu 1} \sim m_{\nu 2}$. The predicted value for the effective Majorana mass $\langle m_{ee} \rangle \simeq 0.0039$ eV in the neutrinoless double beta decay will be within our reach of the future experiments.

The big ansatz is the existence of the X_2 term in the charged lepton sector (1.12). At present, there is no idea on this term. Besides, it seems that the present lepton mass structure is ill matched with the charged lepton mass relation [14]

$$\frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3}. \tag{4.1}$$

The purpose in the early stage of the yukawaon model was to predict the charged lepton mass relation (4.1). The bilinear form (1.2) for the charged lepton mass matrix was indispensable to predict [15] the relation (4.1). If we adopt the present scenario, we must reconsider the origin of the charged lepton mass spectrum. However, in this paper, we do not use the relation (4.1), but only use the observed charged lepton mass values as input values. Therefore the bilinear form such as (1.2) is not necessarily required in this paper. Nevertheless, the formula (4.1) is still attractive. On the other hand, it is also attractive that we can predict 12 observables (2+2+2 lepton and up-quark mass ratios, and 4+2 PMNS mixing parameters) under 4 adjustable parameters $(a_e, a_u, \alpha_u, \zeta_u)$ if once we accept this ansatz (1.12). It is a future task how to understand the existence of X_2 term.

In conclusion, although the form M_ν is, at present, not one which is derived from a rigid theoretical ground, the form will offer a suggestive hint for a unification model of quark and lepton mass matrices.

Acknowledgment

One of the authors (YK) is supported by JSPS (No. 21540266).

Appendix A: R charge assignments

In the present model, as well as in the O(3) model, we construct a model without introducing a yukawaon Y_ν by replacing Y_ν by Y_e. The simple way to guarantee that the yukawaon Y_e couples not only to the charged lepton sector but also to the Dirac neutrino sector is to introduce the following R charge assignment,

$$R(\nu^c) = R(e^c) \equiv r_e, \tag{A.1}$$

$$R(H_u) = R(H_d) = 1. \tag{A.2}$$

The R charge of $(\bar{E}_u E^u)$ is free parameter in the form (2.8). For simplicity, we take

$$R(\bar{E}_u E^u) = R(\Theta_{8+1}) = 1. \tag{A.3}$$

Hereafter, we will denote $R(\bar{E}_u)$ and $R(E^u)$ as $\bar{r}_E$ and $1 - \bar{r}_E$, respectively. Each yukawaon is distinguished from other yukawaons by the R charges. If we define a parameter n as

$$n \equiv 2[R(\bar{Y}_R) - R(\bar{Y}_e)], \tag{A.4}$$

then, we can express the R charges of the other fields from Eq.(2.2) as follows:

$$R(\ell) = r_e + \frac{1}{2}(n-2), \tag{A.5}$$

$$R(\bar{Y}_e) = \frac{1}{2}(4-n) - 2r_e, \tag{A.6}$$

$$R(\bar{Y}_R) = 2 - 2r_e, \tag{A.7}$$

$$R(\bar{Y}_u) = n - 1 + \bar{r}_E, \tag{A.8}$$

$$R(\bar{Y}_d) = \frac{1}{2}(n-2) + \bar{r}_E, \tag{A.9}$$

$$R(u^c) + R(q) = 2 - n - \bar{r}_E, \tag{A.10}$$

$$R(d^c) + R(q) = 2 - \frac{1}{2}n - \bar{r}_E. \tag{A.11}$$

From Eqs.(2.6) and (2.7), we obtain

$$R(\bar{Y}_u) = 2R(\bar{\Phi}_u) + R(E^u), \tag{A.12}$$

$$R(\bar{Y}_R) = R(\bar{\Phi}_u) + R(\bar{Y}_e) + R(E^u), \tag{A.13}$$

respectively. From Eqs.(A.12) and (A.13), we obtain a relation

$$R(\bar{Y}_u) = n - R(E^u) = n - 1 + \bar{r}_E. \tag{A.14}$$

The relation (A.14) leads to

$$R(\bar{Y}_u \Theta^u) = (n-1)R(\bar{E}_u E^u) + R(\bar{E}_u \Theta^u). \tag{A.15}$$

Only when the value n is a positive integer, Eq.(A.15) means that an additional term

$$\frac{\lambda_{0u}}{\Lambda^{2n-1}}\mathrm{Tr}[(\bar{E}_u E^u)^{n-1}\bar{E}\Theta^u], \tag{A.16}$$

can appear in the expression (2.6). Note that if n is not a positive integer, the factor $(\bar{E}_u E^u)^{n-1}$ does not have a physical meaning, because a term with $(E^u)^{-1}$ cannot appear in the superpotential terms. Therefore, the n defined in Eq.(A.4) is allowed only for $n = 1, 2, \cdots$.

As we see in Eqs.(A.5) - (A.11), these R charges are described by four parameters r_e, $R(q)$, $\bar{r}_E$ and n. Therefore, in order to fix these R charge values, we have to assume four constraints for these R charges. On the other hand, the fields $\bar{Y}_e$, $\bar{Y}_R$, $\bar{Y}_u$, $\bar{Y}_d$, $\bar{\Phi}_u$, and $\bar{E}_u$ are gauge singlets, so that they must be distinguished only by R charges. We can choose a suitable parameter set $(n, r_e, r_q, \bar{r}_E)$. Here, let us demonstrate an example of R charge assignments, although it is not the purpose of the present paper to give such an explicit R charge assignment.

For example, we put the following working hypothesis:

$$R(\bar{Y}_\nu) + R(\bar{Y}_e) = 0, \quad R(\bar{Y}_u) + R(\bar{Y}_d) = 0, \tag{A.17}$$

$$R(u^c) + R(d^c) = 0, \quad R(\nu^c) + R(e^c) = 0. \tag{A.18}$$

The constraint (A.17) is an analogy that the Yukawa coupling constants in the standard model do not have R charges. The constraints (A.17) and (A.18) leads to the relation $R(\ell) = R(q) = 1$. Of course, since the yukawaon $\bar{Y}_\nu$ has been replaced by $\bar{Y}_e$ in the present model, the first constraint in Eq.(A.17) reads as $R(\bar{Y}_e) = 0$, and since $R(\nu^c) = R(e^c)$ in the model, the second constraint in Eq.(A.6) reads as $R(e^c) = 0$. Since $R(\bar{Y}_e)$ is given by Eq.(A.7), the requirement $R(\bar{Y}_e) = 0$ together with $R(e^c) = 0$ requires $n = 4$. Thus, the constraints (A.17) and (A.18) fix the parameters $(n, r_e, r_q, \bar{r}_E)$ as

$$n = 4, \quad R(e^c) = 0, \quad R(q) = 1, \quad R(\bar{E}_u) = -2. \tag{A.19}$$

The explicit values of these R values are listed in Table 1. Since the R charges of $\bar{\Phi}_0$, X' and X are still free parameters, we take $R(\bar{\Phi}_0) = \frac{1}{2}$ for simplicity. As we see in Table 1, the fields $\bar{Y}_e$, $\bar{Y}_R$, $\bar{Y}_u$, $\bar{Y}_d$ and $\bar{E}_u$ can safely have different R charges from each other.

Thus, the assumption can lead to plausible R charge values (A.19), so that we consider that the assumption is reasonable. Now we have an additional term,

$$\frac{\lambda_{0u}}{\Lambda^5}\mathrm{Tr}[(\bar{E}_u E^u)^3\bar{E}_u\Theta^u], \tag{A.20}$$

which should be included in M_u given in Eq.(2.6).

Appendix B: Rotation from S_3 into S_2

We define

$$S_3 = \frac{1}{3}\begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad S_2 = \frac{1}{2}\begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{B.1}$$

which are invariant under permutation symmetries S_3 and S_2, respectively. A rotation matrix R which transforms the matrix S_3 into S_2 has been discussed in Ref.[16]. The rotation matrix R is given by

$$R S_3 R^T = S_2, \tag{B.2}$$

where R is defined as follows:

$$R = R_3(-\frac{\pi}{4})\, T\, R_3(\theta)\, A, \tag{B.3}$$

$$R_3(\theta) = \begin{pmatrix} \cos\theta & \sin\theta & 0 \\ -\sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{B.4}$$

$$T = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \tag{B.5}$$

$$A = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{6}} & -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{pmatrix}. \tag{B.6}$$

The matrix A is known as a matrix which diagonalizes the matrix S_3 into

$$A\, S_3\, A^T = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \equiv Z_3, \tag{B.7}$$

and also

$$TA\, S_3\, (TA)^T = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \equiv Z_1. \tag{B.8}$$

The explicit form of R is given by

$$R = \begin{pmatrix} \frac{1}{\sqrt{6}} - \frac{c}{2\sqrt{3}} + \frac{s}{2} & \frac{1}{\sqrt{6}} - \frac{c}{2\sqrt{3}} - \frac{s}{2} & \frac{1}{\sqrt{6}} + \frac{c}{\sqrt{3}} \\ \frac{1}{\sqrt{6}} + \frac{c}{2\sqrt{3}} - \frac{s}{2} & \frac{1}{\sqrt{6}} + \frac{c}{2\sqrt{3}} + \frac{s}{2} & \frac{1}{\sqrt{6}} - \frac{c}{\sqrt{3}} \\ \frac{c}{\sqrt{2}} + \frac{s}{\sqrt{6}} & -\frac{c}{\sqrt{2}} + \frac{s}{\sqrt{6}} & -\frac{2s}{\sqrt{6}} \end{pmatrix}, \tag{B.9}$$

where $s = \sin\theta$ and $c = \cos\theta$.

In the expression (B.9) of R, when we define

$$\begin{aligned} z_1 &= \frac{1}{\sqrt{6}} - \frac{c}{2\sqrt{3}} + \frac{s}{2}, \\ z_2 &= \frac{1}{\sqrt{6}} - \frac{c}{2\sqrt{3}} - \frac{s}{2}, \\ z_3 &= \frac{1}{\sqrt{6}} + \frac{c}{\sqrt{3}}, \end{aligned} \tag{B.10}$$

the rotation matrix R is expressed as follows:

$$R = \begin{pmatrix} z_1 & z_2 & z_3 \\ \sqrt{\frac{2}{3}} - z_1 & \sqrt{\frac{2}{3}} - z_2 & \sqrt{\frac{2}{3}} - z_3 \\ \sqrt{\frac{2}{3}}(z_3 - z_2) & \sqrt{\frac{2}{3}}(z_1 - z_3) & \sqrt{\frac{2}{3}}(z_2 - z_1) \end{pmatrix}. \tag{B.11}$$

Here, z_i $(i = 1, 2, 3)$ satisfies

$$z_1^2 + z_2^2 + z_3^2 = 1, \tag{B.12}$$

and we can choose z_i such as

$$z_1 + z_2 + z_3 = \sqrt{\frac{3}{2}}. \tag{B.13}$$

Suggested from the charged lepton mass relation (4.1), if we choose z_i as

$$z_i = \frac{\sqrt{m_{ei}}}{\sqrt{m_{e1} + m_{e2} + m_{e3}}}, \tag{B.14}$$

where $(m_{e1}, m_{e2}, m_{e3}) = (m_e, m_\mu, m_\tau)$, then, the matrix R satisfies

$$R \begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}. \tag{B.15}$$

Since

$$ZS_3Z = \frac{1}{3} \begin{pmatrix} z_1^2 & z_1 z_2 & z_1 z_3 \\ z_1 z_2 & z_2^2 & z_2 z_3 \\ z_1 z_3 & z_2 z_3 & z_3^2 \end{pmatrix}, \tag{B.16}$$

where

$$Z = \begin{pmatrix} z_1 & 0 & 0 \\ 0 & z_2 & 0 \\ 0 & 0 & z_3 \end{pmatrix}, \tag{B.17}$$

the following relation holds:

$$RZS_3ZR^T = \frac{1}{3}Z_1, \tag{B.18}$$

where Z_1 is defined by Eq.(B.8). However, note that, from Eqs.(B.8) and (B.18), we cannot conclude $RZ = (1/\sqrt{3})TA$.

Thus, it seems the rotation matrix R from S_3 into S_2 is deeply related to the charged lepton mass relation (4.1), but it is not clear why the form S_2 appears in the charge lepton sector. This is still an open question at present.

References

[1] Y. Koide, Phys. Rev. **D 79**, 033009 (2009).

[2] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. **B 147**, 277 (1979).

[3] Y. Koide, Phys. Lett. **B 680**, 76 (2009).

[4] H. Nishiura and Y. Koide, Phys. Rev. **D 83**, 035010 (2011).

[5] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).

[6] H. Fritzsch, Phys. Lett. **73B**, 317 (1978); **85B**, 81 (1979); Nucl. Phys. **B155**, 189 (1979).

[7] B. Pontecorvo, Zh. Eksp. Teor. Fiz. **33**, 549 (1957) and **34**, 247 (1957); Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28**, 870 (1962).

[8] Z.-z. Xing, H. Zhang and S. Zhou, Phys. Rev. **D 77**, 113016 (2008). And also see, H. Fusaoka and Y. Koide, Phys. Rev. **D 57**, 3986 (1998).

[9] Particle Data Group, K. Nakamura, *et al.*, J. Phys. **G 37**, 075021 (2010).

[10] M. Doi, T. Kotani, H. Nishiura, K. Okuda, and E. Takasugi, Phys. Lett. **B103**, 219 (1981) and **B113**, 513 (1982).

[11] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. **B 458**, 79 (1999); Phys. Lett. **B 530**, 167 (2002); Z.-z. Xing, Phys. Lett. **B 533**, 85 (2002); P. F. Harrison and W. G. Scott, Phys. Lett. **B 535**, 163 (2002); Phys. Lett. **B 557**, 76 (2003); E. Ma, Phys. Rev. Lett. **90**, 221802 (2003); C. I. Low and R. R. Volkas, Phys. Rev. **D 68**, 033007 (2003).

[12] K. Abe, *et al.* (T2K Collaboration), Phys. Rev. Lett. **107**, 041801 (2011).

[13] H. De Kerret, a talk at Low Nu, Seoul, Nov. 2011: http://www.dchooz.org/DocDB /0033/003393/003/DCAtLowNu11_Kerret111109_Official.pdf.

[14] Y. Koide, Lett. Nuovo Cimento **34**, 201 (1982); Phys. Lett. **B120**, 161 (1983); Phys. Rev. **D28**, 252 (1983).

[15] Y. Koide, Mod. Phys. Lett. **A5**, 2319 (1990).

[16] Y. Koide and H. Fusaoka, Phys.Rev. **D 66**, 113004 (2002).